

## Recombinant Human Parathyroid Hormone/PTH Protein (aa 32-65, GST Tag)

Catalog Number: PKSH030707

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human Parathyroid Hormone/PTH protein Ser 32-Phe 65, with an N-terminal GST
<b>Calculated MW</b>	31.0 kDa
<b>Observed MW</b>	30 kDa
<b>Accession</b>	P01270
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 96 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.5 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 96 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Parathyroid hormone (PTH); parathormone or parathyrin; is secreted by the chief cells of the parathyroid glands as a polypeptide. PTH elevates calcium level by dissolving the salts in bone and preventing their renal excretion. Parathyroid hormone (PTH) has been proved to play a pivotal role in maintaining myocardial contractility as well as effective natriuresis; and possible pathogenic mechanisms contributing to heart failure secondary to hypocalcemia and hypoparathyroidism. With the increased population of preosteoblastic lineages and the osteoblastic activation; Parathyroid hormone (PTH) drives anabolism in bone. Experiments have recently reported that PTH affects bone cells in a dual pathway - mediating osteoblastic (preosteoblastic) activities or osteocytic synthesis of sclerostin. Defects in PTH are a cause of familial isolated hypoparathyroidism (FIH); also called autosomal dominant hypoparathyroidism or autosomal dominant hypocalcemia. FIH is characterized by hypocalcemia and hyperphosphatemia due to inadequate secretion of parathyroid hormone. Symptoms are seizures; tetany and cramps.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017