Recombinant Mouse CNDP2/CPGL/PEPA Protein (His Tag)

Catalog Number: PKSM041231



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-					
- 10	AC	cri	m		nn
\mathbf{L}			ш	UΙU	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

 Species
 Mouse

 Mol_Mass
 53.8 kDa

 Accession
 O9D1A2

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU per } \mu\text{g of the protein as determined by the LAL method.}$

Storage Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.

Shipping This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel


packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, 10% Glycerol,

pH 8.0.

Reconstitution Not Applicable

Data



Background

Mouse cytosolic non-specific dipeptidase(CNDP2) is a cytoplasm protein which belongs to the peptidase M20A family. CNDP2 has 2 Isoform: Isoform 1 is ubiquitously expressed with higher levels in kidney and liver (at protein level). Isoform 2 is expressed in fetal tissues, it is only expressed in adult liver and placental tissues. CNDP2 hydrolyzes a variety of dipeptides including L-carnosine and has a strong preference for Cys-Gly. It is a cytosolic enzyme that can hydrolyze carnosine to yield 1-histidine and beta-alanine. CNDP2 is highly expressed in the histaminergic neurons in the tuberomammillary nucleus, implying that it may supply histidine to histaminergic neurons for histamine synthesis. It may play a role as tumor suppressor in hepatocellular carcinoma (HCC) cells.

For Research Use Only