Elabscience®

PE/Elab Fluor[®] 594 Anti-Mouse/Human CD11b Antibody[M1/70]

Catalog Number: E-AB-F1081UP

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Reactivity	Human;Mouse	
Host	Rat	
lsotype	Rat lgG2b, κ	
Clone No.	M1/70	
Isotype Control	PE/Elab Fluor [®] 594 Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09843P]	
Conjugation	PE/Elab Fluor [®] 594	
Conjugation Information PE/Elab Fluor [®] 594 is designed to be excited by the blue (488 nm), Green (53 yellow-green (561 nm) lasers and detected using an optical filter centered near		
	(e.g., a 610/20 nm bandpass filter).	
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.	
Applications	Recommended usage	
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the	

reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶ cells in 100 μ L volume].

Data



C57BL/6 murine bone marrow cells are stained with Elab

Fluor[®] Red 780 Anti-Mouse CD45R/B220 Antibody and PE/Elab Fluor[®] 594 Anti-Mouse/Human CD11b Antibody (Left). Bone marrow cells are stained with Elab Fluor[®] Red 780 Anti-Mouse CD45R/B220 Antibody and PE/Elab Fluor[®] 594 Rat IgG2b, κ Isotype Control (Right).

Preparation & Storage		
Storage	Keep as concentrated solution.	
	This product can be stored at 2-8°C for 12 months. exposure to light and do not freeze.	Please protected from prolonged
Shipping	Ice bag	
Antigen Information		
Alternate Names	CD11 antigen-like family member B;CD11b;CR-3 alpha chain;Integrin alpha-M;Itgam; Leukocyte adhesion receptor MO1	
For Research Use Only	/	
Tel: 400-999-2100	Web: w w w .elabscience.cn	Email:techsupport@elabscience.cn

Elabscience®

Uniprot ID Gene ID Background

P05555;P11215

16409,3684

CD11b is a 170 kD glycoprotein also known as αM integrin, Mac-1 α subunit, Mol, CR3, and Ly-40. CD11b is a member of the integrin family, primarily expressed on granulocytes, monocytes/macrophages, dendritic cells, NK cells, and subsets of T and B cells. CD11b non-covalently associates with CD18 (β2 integrin) to form Mac-1. Mac-1 plays an important role in cell-cell interaction by binding its ligands ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4 (CD242), iC3b, and fibrinogen.