Recombinant Human IL-1RAcP/IL-1R3 Protein (His Tag)

Catalog Number: PKSH031857

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	HEK293 Cells-derived Human IL-1RAcP/IL-1R3 protein Met 1-Glu 359, with an C-
	terminal His
Calculated MW	40.7 kDa
Observed MW	50-55 kDa
Accession	NP_002173.1
Bio-activity	Not validated for activity
Properties	
Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

Web:www.elabscience.com

Elabscience®

Interleukin-1 receptor accessory protein (IL-1RAcP) also known as Interleukin-1 receptor member 3 (IL-1R3) is a a cytokine receptor which binds interleukin 1. The IL-1 receptor accessory protein (IL1RAP) is a transmembrane protein that interacts with IL-1R and is required for IL-1 signal transduction. Interleukin 1 induces synthesis of acute phase and proinflammatory proteins during infection, tissue damage, or stress, by forming a complex at the cell membrane with an interleukin 1 receptor and an accessory protein. IL-1RAcP/IL-1R3 is a necessary part of the interleukin 1 receptor complex which initiates signalling events that result in the activation of interleukin 1-responsive genes. Alternative splicing of this gene results in two transcript variants encoding two different isoforms, one membrane-bound and one soluble. The ratio of soluble to membrane-bound forms increases during acute-phase induction or stress. IL-1RAcP/IL-1R3 mediates interleukin-1-dependent activation of NF-kappa-B. Isoform 1 is part of the membrane-bound form of the IL-1 receptor. Signaling involves formation of a ternary complex containing IL1R1, TOLLIP, MYD88, and IRAK1 or IRAK2. Isoform 2 modulates the response to interleukins by associating with soluble IL1R1 and enhancing interleukin-binding to the decoy receptor.