

## Recombinant Human DSC2/Desmocollin-2 Protein (His Tag)

**Catalog Number:** PKSH031417

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human DSC2/Desmocollin-2 protein Met 1-Arg 684, with an C-terminal His
<b>Calculated MW</b>	75.0 kDa
<b>Observed MW</b>	85&100 kDa
<b>Accession</b>	Q02487-1
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 75 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 75 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

DSC2 is a calcium-dependent glycoprotein that is a member of the desmocollin subfamily of the cadherin superfamily. Like other desmocollins, murine DSC2 has two products, Dsc2a and Dsc2b, produced by alternative splicing of a 46 bp exon which encodes 11 COOH-terminal aa followed by an in-frame stop codon. These desmosomal family members, along with the desmogleins, are found primarily in epithelial cells where they constitute the adhesive proteins of the desmosome cell-cell junction and are required for cell adhesion and desmosome formation. The desmosomal family members are arranged in two clusters on chromosome 18, occupying less than 650 kb combined. Mutations in DSC2 are associated with arrhythmogenic right ventricular dysplasia-11. DSC2 is involved in the interaction of plaque proteins and intermediate filaments mediating cell-cell adhesion. DSC2 may contribute to epidermal cell positioning by mediating differential adhesiveness between cells that express different isoforms.