

Recombinant Human DSC2/Desmocollin-2 Protein (His Tag)

Catalog Number: PKSH031417

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human DSC2/Desmocollin-2 protein Met 1-Arg 684, with an C-terminal His
Calculated MW	75.0 kDa
Observed MW	85&100 kDa
Accession	Q02487-1
Bio-activity	Not validated for activity

Properties

Purity	> 75 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 75 % as determined by reducing SDS-PAGE.

Background

DSC2 is a calcium-dependent glycoprotein that is a member of the desmocollin subfamily of the cadherin superfamily. Like other desmocollins, murine DSC2 has two products, Dsc2a and Dsc2b, produced by alternative splicing of a 46 bp exon which encodes 11 COOH-terminal aa followed by an in-frame stop codon. These desmosomal family members, along with the desmogleins, are found primarily in epithelial cells where they constitute the adhesive proteins of the desmosome cell-cell junction and are required for cell adhesion and desmosome formation. The desmosomal family members are arranged in two clusters on chromosome 18, occupying less than 650 kb combined. Mutations in DSC2 are associated with arrhythmogenic right ventricular dysplasia-11. DSC2 is involved in the interaction of plaque proteins and intermediate filaments mediating cell-cell adhesion. DSC2 may contribute to epidermal cell positioning by mediating differential adhesiveness between cells that express different isoforms.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017