# **Elabscience**®

## Human ADAM10 Antibody Pair Set

Catalog No.E-KAB-0259ApplicationsELISASynonymsAD10, CD156c, CDw156, HsT18717, MADM, kuz, ADAM metallopeptidase domain 10

#### Kit components & Storage

Title	Specifications	Storage
Human ADAM10 Capture Antibody	1 vial, 100 µ g	Store at $-20^{\circ}$ C for one year.
		Avoid freeze / thaw cycles.
Human ADAM10 Detection Antibody	1 vial, 50 μL	Store at $-20^{\circ}$ C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0259)	
		Human ADAM10 Capture Antibody	Human ADAM10 Detection Antibody (Biotin)
Immunogen	Immunogen	Recombinant Human ADAM10	Recombinant Human ADAM10 protein
Information		protein	
	Swissprot	O14672	
Product details Reactivity		Human	Human
	Host	Mouse	Mouse
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Protein A or G	Protein A or G
Specificity		Detects Human ADAM10 in ELISAs.	

For Research Use Only

# **Elabscience**®

### Applications

Human ADAM10 Sandwich ELISA Assay:

	Concentration/Dilution 0.5-4µg/mL	Human ADAM10 Capture	
	0.5-4µg/mL	Human ADAM10 Capture	
Capture			
		Antibody	
ELISA 1 Detection	1:1000-1:10000	Human ADAM10 Detection Antibody (Biotin)	0.01 0.01 0.01 0.01 0.01 0.01 0.01 0.00 0.00 0.00 0.000 0.00000 0.000000

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

### Background

Members of the ADAM family are cell surface proteins with a unique structure possessing both potential adhesion and protease domains. This gene encodes and ADAM family member that cleaves many proteins including TNF-alpha and E-cadherin. Cleaves the membrane-bound precursor of TNF-alpha at '76-Ala-|-Val-77' to its mature soluble form. Responsible for the proteolytical release of soluble JAM3 from endothelial cells surface. Responsible for the proteolytic release of several other cell-surface proteins, including heparin-binding epidermal growth-like factor, ephrin-A2 and for constitutive and regulated alpha-secretase cleavage of amyloid precursor protein (APP).