Recombinant Mouse PRLR/Prolactin Receptor Protein (Fc Tag)



Catalog Number: PKSM041287

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Prolactin receptor;PRL-R;Prlr;Prolactin R;PRLR

Species Mouse

Expression Host HEK293 Cells
Sequence Gln20-Asp229

AccessionQ08501Calculated Molecular Weight51.7 kDaObserved molecular weight62-88 kDaTagC-Fc

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

The prolactin receptor (PRLR) is a member of the class I cytokine/lactogen receptor family which mediates the diverse cellular actions of prolactin in several tissues. PRLRs are expressed in normal and neoplastic human breast tissue, and in most breast cancer cells. PRLR contains an extracellular region that binds prolactin, a transmembrane region, and a cytoplasmatic region required for the activation of the Jak2–Stat5 signal transduction pathway by Prl which is essential for transcriptional activation of all known prolactin regulated genes. PRLRs have also been observed in ovarian follicular cells of mice, pigs, sheep, deer, and humans, as well as in luteal tissue in cow and horse ovaries. Furthermore, PRLR knockout mice exhibit failure of embryonic implantation, reduced number of mature oocytes, and low fertilization rates. Knockout females also display a reduced number of primary follicles.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com