Elabscience®

LMAN2L Polyclonal Antibody

catalog number: E-AB-92075

Note: Centrifuge before opening to ensure complete recovery of vial contents.

1:50-1:200

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human LMAN2L
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000

Data

IF



Western blot analysis of extracts of HeLa cells using LMAN2L Polyclonal Antibody at 1:1000 dilution.





Immunofluorescence analysis of C6 cells using LMAN2L Polyclonal Antibody at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using LMAN2L Polyclonal Antibody at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining. Immunofluorescence analysis of U2OS cells using LMAN2L Polyclonal Antibody at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.

Preparation & Storage

Storage Shipping Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:<u>techsupport@elabscience.com</u> Fax: 1-832-243-6017

Elabscience®

This gene encodes a protein belonging to the L-type lectin group of type 1 membrane proteins, which function in the mammalian early secretory pathway. These proteins contain luminal carbohydrate recognition domains, which display homology to leguminous lectins. Unlike other proteins of the group, which cycle in the early secretory pathway and are predominantly associated with post endoplasmic reticulum membranes, the protein encoded by this gene is a non-cycling resident protein of the ER, where it functions as a cargo receptor for glycoproteins. It is proposed to regulate exchange of folded proteins for transport to the Golgi and exchange of misfolded glycoproteins for transport to the ubiquitin-proteasome pathway. [provided by RefSeq, Apr 2016]

For Research Use Only