

## Recombinant Human PACSIN2/Syndapin-2 Protein (His Tag)

Catalog Number: PKSH032846

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human PACSIN2;Syndapin-2 protein Met 1-Gln486, with an C-terminal His
<b>Calculated MW</b>	56.7 kDa
<b>Observed MW</b>	88 kDa
<b>Accession</b>	Q9UNF0
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
	Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Protein Kinase C and Casein Kinase Substrate in Neurons Protein 2 (PACSIN2) is a member of the PACSIN family. PACSIN2 is localized to the plasma membrane via its coiled-coil domain. PACSIN2 is widely expressed and contains one FCH domain and one SH3 domain. PACSIN2 forms homo- and hetero-aggregates with other PACSINs. PACSIN2 may play a role in vesicle formation and transport. In addition, PACSIN2 is involved in linking the actin cytoskeleton with vesicle formation by regulating tubulin polymerization.

### For Research Use Only