
Recombinant Rat GSTM2 protein (His Tag)

Catalog Number: PDER100223

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Species Source	Rat E.coli-derived Rat GSTM2 protein Pro2-Lys218, with an N-terminal His
Coloulated MW	
Calculated MW	23.8 kDa
Observed MW	30 kDa
Accession	P08010
Bio-activity	Not validated for activity
Properties	
Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with 5% Trehalose and 5%
	Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of
	0.5 mg/mL. Concentration is measured by UV-Vis.



> 95 % as determined by reducing SDS-PAGE.

Background

Data

Glutathione S-transferase Mu 2, also known as GST class-mu 2, GSTM2-2 and GSTM2, is a cytoplasm protein which belongs to the GST superfamily and Mu family. GSTM2 / GST4 contains one ,GST C-terminal domain and one& nbsp,GST N-terminal domain. The glutathione S-transferases (GSTs) are a multigene family of enzymes largely involved in the detoxification of chemicals. Eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. ,Butyrate, an important luminal component produced from fermentation of dietary fibers, is an efficient inducer of GSTs and especially of GSTM2. Butyrate may act chemoprotectively by increasing detoxification capabilities in the colon mucosa.

Web:www.elabscience.com