

# ERK 1/2 Polyclonal Antibody

catalog number: D-AB-10257L

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description


<b>Reactivity</b>	Mouse
<b>Immunogen</b>	Recombinant Rat ERK 1/2 protein expressed by E.coli
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Antigen Affinity Purification
<b>Conjugation</b>	Unconjugated
<b>buffer</b>	PBS with 0.05% proclin 300, 1% protective protein and 50% glycerol,pH7.4

## Applications

## Recommended Dilution

<b>WB</b>	1:500-1:1000
-----------	--------------

## Data



Western blot with ERK 1/2 Polyclonal Antibody at dilution  
of 1:1000.lane 1:NIH/3T3 whole cell lysate

**Observed-MV:43 kDa**

**Calculated-MV:43 kDa**

## Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

## Background

The protein encoded by this gene is a member of the MAP kinase family. MAP kinases,also known as extracellular signal-regulated kinases (ERKs),act in a signaling cascade that regulates various cellular processes such as proliferation,differentiation,and cell cycle progression in response to a variety of extracellular signals. This kinase is activated by upstream kinases,resulting in its translocation to the nucleus where it phosphorylates nuclear targets. Alternatively spliced transcript variants encoding different protein isoforms have been described,This gene encodes a member of the MAP kinase family. MAP kinases,also known as extracellular signal-regulated kinases (ERKs),act as an integration point for multiple biochemical signals,and are involved in a wide variety of cellular processes such as proliferation, differentiation,transcription regulation and development. The activation of this kinase requires its phosphorylation by upstream kinases. Upon activation,this kinase translocates to the nucleus of the stimulated cells,where it phosphorylates nuclear targets. One study also suggests that this protein acts as a transcriptional repressor independent of its kinase activity. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. Two alternatively spliced transcript variants encoding the same protein,but differing in the UTRs,have been reported for this gene.

## For Research Use Only