

## Recombinant Phospho-NF-κB p65 (Ser536) Monoclonal Antibody

catalog number: **AN301332L**

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

### Description


<b>Reactivity</b>	Human;Mouse;Rat;Chicken
<b>Immunogen</b>	A synthetic peptide corresponding to residues around (Ser536) of Human Phospho-NF-κB p65
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG,κ
<b>Clone</b>	B1095
<b>Purification</b>	Protein A
<b>Buffer</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

### Applications

### Recommended Dilution

<b>WB</b>	1:2000-1:10000
<b>IF</b>	1:200-1:1000
<b>ELISA</b>	1:5000-1:20000
<b>IP</b>	1:50-1:200

### Data



Western Blot with Recombinant Phospho-NF-κB p65 (Ser536) Monoclonal Antibody at dilution of 1:1000 dilution. Lane A: NIH-3T3 whole cell lysate.

**Observed-MW:60 kDa**

**Calculated-MW:60 kDa**

### Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	Ice bag

### Background

NF-kappa-B is a ubiquitous transcription factor involved in several biological processes. It is held in the cytoplasm in an inactive state by specific inhibitors. Upon degradation of the inhibitor, NF-kappa-B moves to the nucleus and activates transcription of specific genes. NF-kappa-B is composed of NFKB1 or NFKB2 bound to either REL, RELA, or RELB. The most abundant form of NF-kappa-B is NFKB1 complexed with the product of this gene, RELA. Four transcript variants encoding different isoforms have been found for this gene.

### For Research Use Only