

Purified Anti-Human CD314 Antibody[1D11]

catalog number: E-AB-F1383A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|---------------------|---|
| Reactivity | Human |
| Immunogen | Recombinant Human CD314 protein |
| Host | Mouse |
| Isotype | Mouse IgG1, κ |
| Clone | 1D11 |
| Purification | >98%, Protein A/G purified |
| Buffer | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling. |

Applications

Recommended Dilution

| | |
|------------|--|
| FCM | 2 µg/mL (0.5×10 ⁶ -1×10 ⁶ cells) |
|------------|--|

Data



Verified Samples in FCM: Human peripheral blood lymphocytes

Preparation & Storage

| | |
|----------------|--|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
|----------------|--|

Background

Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. The NKG2 gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed in NK cells. This gene encodes a member of the NKG2 family. The encoded transmembrane protein is characterized by a type II membrane orientation (has an extracellular C terminus) and the presence of a C-type lectin domain. It binds to a diverse family of ligands that include MHC class I chain-related A and B proteins and UL-16 binding proteins, where ligand-receptor interactions can result in the activation of NK and T cells. The surface expression of these ligands is important for the recognition of stressed cells by the immune system, and thus this protein and its ligands are therapeutic targets for the treatment of immune diseases and cancers. Read-through transcription exists between this gene and the upstream KLRC4 (killer cell lectin-like receptor subfamily C, member 4) family member in the same cluster.

For Research Use Only