

Recombinant Phospho-SMAD2 (Ser465, 467) Monoclonal Antibody

catalog number: AN300146L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Human
Immunogen	A synthetic peptide corresponding to the residues around (Ser465, 467) of Human Phospho-SMAD2
Host	Rabbit
Isotype	IgG
Clone	4B12
Purification	Protein A
Buffer	10 mM sodium HEPES, 150 mM NaCl, 100 µg/mL protein protectant, 50% glycerol, pH 7.5


Applications

Recommended Dilution

WB

1:500-1:2000



Data



Western blot analysis of extracts from serum-starved HeLa, untreated (line A) or treated with TGF beta 1 (5 ng/mL, 120min; +) (line B), using Phospho-SMAD2 (Ser465, 467) Monoclonal Antibody at 1:1000 dilution (upper) or Anti-SMAD2 Monoclonal Antibody at 1:100000 dilution (lower).

Observed-MW:60 kDa

Calculated-MW:52 kDa



Western blot analysis of extracts from serum-starved HeLa, untreated (line A); treated with TGFβ1 (5 ng/mL, 120 min), without peptide (line B) or antigen-specific phosphopeptide (line C) or antigen-specific peptide (line D) using Phospho-SMAD2 (Ser465, 467) Monoclonal Antibody at 1:1000 dilution.

Observed-MW:60 kDa

Calculated-MW:52 kDa

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.1

Western blot analysis of extracts from serum-starved Hela, untreated (line A); treated with TGFβ1 (5 ng/mL, 120min; +); treated with TGFβ1 and λ-phosphatase (line C) using Phospho-SMAD2 (Ser465, 467) Monoclonal Antibody at 1:1000 dilution.

Observed-MW:60 kDa

Calculated-MW:52 kDa

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the *Drosophila* gene 'mothers against decapentaplegic' (Mad) and the *C. elegans* gene *Sma*. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 2 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants have been observed for this gene.

For Research Use Only