Elabscience®

Recombinant S100A8 Monoclonal Antibody

catalog number: AN300036P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

ъ	, , ,	
Description		
Reactivity	Human	
Immunogen	Recombinant Human S1	00A8 Protein
Host	Rabbit	
Isotype	IgG	
Clone	8D8	
Purification	Protein A	
Buffer	0.2 µm filtered solution i	n PBS
Applications	Recommended Dilu	ution
WB	1:500-1:2000	
IP	5-10 μ L/mg of lysate	
Data		
KDa A 95 72 55 34 26 17	←── IgG heavy chain	KDa A 95 72 55 43 34 26 17
1100	< S100A8	\rightarrow
Immunoprecipitation	analysis using 4 µL anti-S100A8	Western Blot with \$100A8 Monoclonal Antibody at dilution

Immunoprecipitation analysis using 4 µL anti-S100A8 Monoclonal Antibody and 60 µg of Immunomagnetic beads Protein A/G. Western blot was performed from the immunoprecipitate using S100A8 Monoclonal Antibody at a dilution of 1:100. Lane A:0.5 mg THP-1 Whole Cell Lysate

Observed-MW:11 kDa Calculated-MW:11 kDa

Western Blot with S100A8 Monoclonal Antibody at dilution of 1:500. Lane A: THP-1 Whole Cell Lysate, Lysates/proteins at 30 µg per lane. **Observed-MW:11 kDa**

Calculated-MW:11 kDa

Calculated-MW:11 kDa	
Preparation & Storage	
Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag
Background	

Elabscience®

S100A8/S100A9 HeterodimerS100A8 (also MRP8 and calgranulin A) is a 10 kDa member of the S100 family, EF-hand superfamily of Ca-binding proteins. It is produced by neutrophils and monocytes, and forms Ca2+-dependent heterodimer/heterotetramer complexes (termed calprotectin) with S100A9. It functions both intracellularly and extracellularly, where it binds to RAGE and CD36. Human S100A8 is 93 amino acids (aa) in length. It contains two EF-hand motifs (aa 12 - 47 and 46 - 81) and one high-affinity Ca2+-binding site (aa 59 - 70). There may be one splice form that shows a 15 aa substitution for the C-terminal 14 amino acids. Although mouse S100A8 is cleaved by MMP-2 after Asn21, it is unclear if human S100A8 is susceptible. Full-length human S100A8 is 57% and 74% aa identical to mouse and canine S100A8, respectively.