

## Recombinant Human NT protein (His Tag)

**Catalog Number:** PDEH100801

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human NT protein Ser24-Leu148, with an N-terminal His
<b>Calculated MW</b>	13.6 kDa
<b>Observed MW</b>	17 kDa
<b>Accession</b>	P30990
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 10 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

DPEP2 (MBD-2) belongs to the membrane-bound dipeptidase family. There are three members of this family as membrane-bound dipeptidase-1 (MBD-1), membrane-bound dipeptidase-2 (MBD-2) and membrane-bound dipeptidase-3 (MBD-3). MBD-2 is expressed at highest levels in lung, heart, and testis and at some what lower levels in spleen. MBD-2 is membrane-bound through a glycosylphosphatidylinositol linkage and probably is a metalloprotease which hydrolyzes leukotriene D4 (LTD4) into leukotriene E4 (LTE4). It is generally recognized that rapid cleavage of LTD4 is important in inactivating the broncho- and vaso-constrictive effects of cysteinyl LTs in asthmatic and inflammatory processes.

### For Research Use Only