Elab Fluor[®] 488-conjugated beta Tubulin Monoclonal Antibody

catalog number: AN00293L




Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Human;Mouse;Rat;Zebrafish |
| Immunogen | Recombinant human TUBB protein by E. Coli |
| Host | Mouse |
| Isotype | IgG |
| Clone | 2E8-2 |
| Purification | Protein A/G Purification |
| Conjugation | Elab Fluor [®] 488(Ex/Em:493nm/519nm) |
| buffer | PBS with 0.05% proclin 300, 1% protective protein and 50% glycerol,pH7.4 |
| Applications | Recommended Dilution |

Арр IF Recommended Dilution 1:100-1:200





Immunofluorescence analysis of Hela cells using Elab Fluor

[®] 488-conjugated beta Tubulin Monoclonal Antibody at dilution of 1:100

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Protected from |
| | prolonged exposure to light. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the |
| | temperature recommended. |
| | temperature recommended. |

Background

In eukaryotes there are six members of the tubulin superfamily. α - and β -tubulins polymerize into microtubules, a major component of the eukaryotic cytoskeleton. Microtubules function in many essential cellular processes, including mitosi s. Both α and β tubulins have a mass of around 50 kDa. Tubulin was long thought to be specific to eukaryotes. Beta-tubulin gene is commonly used as a loading control for western blot.

For Research Use Only