



The professional cell culture empowers a healthier world

Retinoic Acid (ATRA) Solution (100 mM)

Cat. No.: PB180614

Size: 1mL

General Information

Product Form Liquid

Solvent

Concentration

Storage

Whether to avoid light

DMSO (100%)

100 mmol/L

-5~-20°C

Shading light

Shipping I lee bag

Expiration date 12 months

Background

Vitamin A acid is a metabolic intermediate of vitamin A. It is also a natural ligand for the Retinoic Acid Receptor (RAR) and the Retinoid X Receptor (RXR), which can induce cell differentiation and apoptosis. These receptors are essential for cell growth, differentiation, and organ formation, as well as for epithelial metabolism and bone growth.

Notes

- 1. This product was sterilized by 0.1 μm filtration and can be used directly after melting.
- 2. It is necessary to pay attention to the aseptic operation and avoid the contamination.
- 3. Before using, the product should be thawed at 2-8°C and shaken thoroughly; repeated freeze-thaw cycles are not advised.
- 4. If precipitation happens after thawing, the contents can be resuspended by pipetting or vortex mixing. After incubating the solution at 37°C for 20 to 30 minutes or letting it stand at room temperature for about an hour, check to see if the precipitate dissolves as intended. If the product dissolves completely, it can be used as usual.
- 5. This product is a concentrated solution and should be diluted prior to use as required.
- 6. The product should be used within a month if stored regularly at 2-8°C. Keep in a frozen state at -5~-20°C for extended storage. Long-term storage at room temperature or between 2-8°C is not recommended. When lesser amounts are required, aliquoting is advised to prevent repeated freeze-thaw cycles.
- 7. This product is for research use only.
- 8. Harmful if swallowed, cause skin irritation, may cause damage to fertility or fetus, extremely toxic to aquatic organisms with long-term persistent effects, please take relevant precautions when handling.

Rev V1.0