Recombinant Swine IL-4 protein(His Tag)

Catalog Number: PKSS000004



Note: Centrifuge before opening to ensure complete recovery of vial contents.

DAG	crin	non
Des	CIID	$\mathbf{u}\mathbf{v}\mathbf{n}$

 Species
 Porcine

 Mol_Mass
 13.5 kDa

 Accession
 Q04745

Bio-activity Measure by its ability to induce proliferation in TF-1 cells. The ED₅₀ for this effect is

<0.6 ng/mL.

Properties

Purity > 98 % as determined by reducing SDS-PAGE.

Endotoxin < 0.1 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

Interleukin-4 (IL-4) is a pleiotropic cytokine that regulates diverse T and B cell responses including cell proliferation; survival and gene expression. IL-4 is produced by mast cells; T cells; and bone marrow stromal cells. IL-4 regulates the differentiation of naive CD4+ T cells into helper Th2 cells; characterized by their cytokine-secretion profile that includes secretion of IL-4; IL-5; IL-6; IL-10; and IL-13; which favor a humoral immune response. Another dominant function of IL-4 is the regulation of immunoglobulin class switching to the IgGl and IgE isotypes. Excessive IL-4 production by Th2 cells has been associated with elevated IgE production and allergic response.

For Research Use Only