

Recombinant Human IL-1 beta/IL1B Protein (pro form, His Tag)

Catalog Number: PKSH031840

Note: Centrifuge before opening to ensure complete recovery of vial contents.



Description

Species	Human
Source	E.coli-derived Human IL-1 protein Met 1-Ser 269(Glu6Lys), with an N-terminal His
Calculated MW	32.3 kDa
Observed MW	34 kDa
Accession	NP_000567.1
Bio-activity	Immobilized human IL1B at 10 µg/mL (100 µl/well) can bind human IL1R1. The EC ₅₀ of human IL1R1 is 0.198 µg/mL.

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



Immobilized human IL1B at 10 µg/mL (100 µl/well) can bind human IL1R1. The EC₅₀ of human IL1R1 is 0.198 µg/mL.

Background

For Research Use Only

Interleukin-1 beta (IL1 beta or IL1B) also known as catabolin, is a member of the interleukin 1 cytokine family. IL1 is a pleiotropic cytokine. It is involved in the inflammatory response, cell growth, and tissue repair in the cortex. The IL1 superfamily consists of three members, IL1A (IL1 alpha), IL1B (IL1 beta), and IL1 receptor antagonist (IL1Ra). In clinical, it has been reported that Interleukin (IL)-1 may influence Th1 / Th2 immune responsiveness and has been implicated in the establishment of successful pregnancy. Proinflammatory interleukin (IL)-1 gene polymorphisms associated with high levels of IL-1beta activity increase the risk for hypochlorhydria and distal gastric carcinoma. IL1B polymorphisms may be involved in susceptibility to SSc. Moreover, the IL2-384-G allele may be a marker for the limited phenotype of systemic sclerosis (SSc).