

A Reliable Research Partner in Life Science and Medicine

# APC Anti-Human CD22 Antibody[HIB22]

Catalog Number: E-AB-F1321E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

## Description

Reactivity Human Host Mouse

**Isotype** Mouse IgG1, κ

Clone No. HIB22

Isotype Control APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]

Conjugation APC

**Conjugation Information** APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


## Applications

## Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### Data



Human peripheral blood lymphocytes are stained with FITC Anti-Human CD19 Antibody and APC Anti-Human CD22 Antibody (Left). Lymphocytes are stained with FITC Anti-Human CD19 Antibody and APC Mouse IgG1, κ Isotype Control (Right).

# **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

# **Antigen Information**

Alternate Names Siglec2;B-cell receptor CD22;B-lymphocyte cell adhesion molecule;BL-CAM;CD22;

Cd22;Lyb-8;Sialic acid-binding lg-like lectin 2;Siglec-2;T-cell surface antigen Leu-14

Uniprot ID P20273

# For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com

## **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

Gene ID Background 933

CD22 is a 130 kD type I transmembrane glycoprotein also known as Siglec-2 and BL-CAM. It is a member of the immunoglobulin superfamily (sialoadhesion subgroup). CD22 is expressed in the cytoplasm of pro-B and pre-B cells, and on the surface of mature B and activated B cells, but not on plasma cells. CD22 is present in the B cell receptor complex and associates with SHP-1, Syk, Lck, Lyn, and phospholipase Cγ1. A primary function of CD22 is thought to be in limiting antigen receptor signaling by modulating B cell activation threshold. CD22 has been shown to bind to CD45RO and CD75, although the natural ligands for this molecule remain controversial.

Fax: 1-832-243-6017