A Reliable Research Partner in Life Science and Medicine

Recombinant Human Cytochrome C/CYCS Protein (His Tag)

Catalog Number: PKSH032336

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source E.coli-derived Human Cytochrome C;CYCS protein Gly2-Glu105, with an C-terminal

His

Calculated MW12.8 kDaObserved MW16 kDaAccessionP99999

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Concentration Subject to label value.

Endotoxin $< 1.0 \text{ EU} \text{ per } \mu\text{g} \text{ of the protein as determined by the LAL method.}$

Storage Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.


Shipping This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel

packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 20mM PB, 10% Trehalose, 200mM NaCl,

50% Glycerol, 0.05% Tween 80, pH7.0.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Cytochrome C (CYCS) is a small heme protein that belongs to the cytochrome c family. It is found loosely associated with the inner membrane of the mitochondrion. Cytochrome C is a highly soluble protein that functions as a central component of the electron transport chain in mitochondria. CYCS transfers electrons between Complexes III (Coenzyme Q - Cyt C reductase) and IV (Cyt C oxidase). CYCS plays a role in apoptosis. Suppression of the anti-apoptotic members or activation of the pro-apoptotic members of the Bcl-2 family leads to altered mitochondrial membrane permeability resulting in release of cytochrome c into the cytosol. Binding of Cytochrome C to Apaf-1 triggers the activation of caspase-9, which then accelerates apoptosis by activating other caspases.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017