

Recombinant Human Alkaline Phosphatase/ALPL Protein (His Tag)

Catalog Number: PKSH031604

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human Alkaline Phosphatase/ALPL protein Met 1-Ser 502, with an C-terminal His
Calculated MW	55.0 kDa
Observed MW	65 kDa
Accession	NP_000469.3
Bio-activity	Measured by its ability to cleave a fluorogenic substrate, 4-Methylumbelliferyl phosphate (4-MUP). The specific activity is > 50, 000 pmoles/min/μg.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 25mM Tris, 0.15M NaCl, pH 7.5 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Alkaline phosphatase (ALPL) is a hydrolase enzyme responsible for removing phosphate groups from many types of molecules, including nucleotides, proteins, and alkaloids. The process of removing the phosphate group is called dephosphorylation. As the name suggests, alkaline phosphatases are most effective in an alkaline environment. It is sometimes used synonymously as basic phosphatase. Alkaline phosphatases (APs) are ubiquitous in many species, from bacteria to human. Four genes encode AP isoenzymes in humans and rodents. Three AP genes are expressed in a tissue-specific manner (i.e., placental, embryonic, and intestinal AP isoenzymes). Expression of the fourth AP gene is nonspecific to a single tissue and is especially abundant in bone, liver, and kidney. This isoenzyme is also called tissue-nonspecific alkaline phosphatase (TNAP). The enzyme tissue non-specific alkaline phosphatase (TNAP) belongs to the ectophosphatase family. TNAP is present in large amounts in bone in which it plays a role in mineralization.