Endoglin/CD105 Polyclonal Antibody(Capture/Detector)


catalog number: AN000260P



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Immunogen	Recombinant Human Endoglin/CD105 protein expressed by E.coli
Host	Rabbit
Isotype	Rabbit IgG
Purification	Antigen Affinity Purification
Conjugation	Unconjugated
buffer	Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.
Applications	Recommended Dilution
ELISA Capture	2-8 µg/mL
ELISA Detector	$0.1-0.4 \ \mu g/mL$

Data



Sandwich ELISA-Recombinant Human Endoglin/CD105 protein standard curve.Background subtracted standard curve using Endoglin/CD105 antibody(AN000260P) (Capture),Endoglin/CD105 antibody(AN000260P)(Detector) in sandwich ELISA.The reference range value for Recombinant Human Endoglin/CD105 protein is 0.15625-10 ng/mL.

Preparation & Storage	
Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /
	thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the
	temperature recommended.

Background

Vascular endothelium glycoprotein that plays an important role in the regulation of angiogenesis.Required for normal structure and integrity of adult vasculature.Regulates the migration of vascular endothelial cells.Required for normal extraembryonic angiogenesis and for embryonic heart development.May regulate endothelial cell shape changes in response to blood flow,which drive vascular remodeling and establishment of normal vascular morphology during angiogenesis.May play a critical role in the binding of endothelial cells to integrins and/or other RGD receptors.Acts as TGF-beta coreceptor and is involved in the TGF-beta/BMP signaling cascade that ultimately leads to the activation of SMAD transcription factors.Required for GDF2/BMP9 signaling through SMAD1 in endothelial cells and modulates TGFB1 signaling through SMAD3.

For Research Use Only