

CD15 Monoclonal Antibody

catalog number: E-AB-22100

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Immunogen	Synthetic Peptide
Host	Mouse
Isotype	IgG
Clone	6B2
Purification	Protein A purification
Conjugation	Unconjugated
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein protectant and 50% glycerol.

Applications


Recommended Dilution

IHC	1:100-1:300
IF	1:50-200

Data



Immunohistochemistry of paraffin-embedded Human lung cancer tissue using CD15 Monoclonal Antibody at dilution of 1:200.



Immunofluorescence analysis of Human liver cancer tissue using CD15 Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

The product of this gene transfers fucose to N-acetylglucosamine polysaccharides to generate fucosylated carbohydrate structures. It catalyzes the synthesis of the non-sialylated antigen, Lewis x (CD15).

FUT4 (Fucosyltransferase 4) is a Protein Coding gene. Diseases associated with FUT4 include Liver Lymphoma and Colon Adenocarcinoma. Among its related pathways are Mannose type O-glycan biosynthesis and Wnt / Hedgehog / Notch. GO annotations related to this gene include fucosyltransferase activity and alpha-(1->3)-fucosyltransferase activity. An important paralog of this gene is FUT5.

For Research Use Only