

Recombinant IL20RA/IL-20RA Monoclonal Antibody

catalog number: AN300332P

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Reactivity	Human
Immunogen	Recombinant Human IL20RA/IL-20RA protein
Host	Rabbit
Isotype	IgG
Clone	1F15
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS

Applications Recommended Dilution

IHC-P	1:100-1:500
--------------	-------------

Data



Immunohistochemistry of paraffin-embedded human skin using IL20RA/IL-20RA Monoclonal Antibody at dilution of 1:200. Positive staining was localized to stratified squamous epithelium.

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

Interleukin 20 receptor, alpha subunit (IL20RA/IL-20RA) also known as IL-20 receptor subunit alpha, Cytokine receptor class-II member 8, interleukin-20 receptor I, and interleukin-20 receptor subunit alpha, is a subunit for the interleukin-20 receptor. IL20RA/IL-20RA belongs to the type II cytokine receptor family. This cytokine is a receptor for interleukin 20 (IL20), a cytokine that may be involved in the epidermal function. The receptor of IL20 is a heterodimeric receptor complex consisting of this protein and interleukin 20 receptor beta (IL20B). IL20RA forms heterodimer with IL20RB and the complex serves as a receptor for IL19, IL20, and IL24. All three are capable of signaling through the IL20RA/IL-20RB complex. The ligand binding to receptor B creating a high-affinity binding site for receptor A which is recruited to complete the complex. Besides, IL20RA also forms a heterodimer with the unique and specific receptor IL10RB and functions as the receptor for IL26. IL20RA is widely expressed with the highest levels in skin, testis, and brain. The expression of both IL20RA and IL20RB is found to be upregulated in psoriatic skin lesions on keratinocytes.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.0