

A Reliable Research Partner in Life Science and Medicine

PerCP/Cyanine5.5 Anti-Mouse CD49d Antibody[R1-2]

Catalog Number: AN00422J

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

lsotype Rat lgG2b, κ

Clone No. R1-2

Isotype Control PerCP/Cyanine5.5 Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09842J]

Conjugation PerCP/Cyanine 5.5

Conjugation Information PerCP/Cyanine5.5 is designed to be excited by the blue laser (488 nm) and detected

using an optical filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine splenocytes cells with PerCP/Cyanine5.5 Anti-Mouse CD49d Antibody[R1-2] (filled gray histogram) or PerCP/Cyanine5.5 Rat IgG2b, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names α4 integrin;VLA-4 α chain;integrin α4;ITGA4

 Uniprot ID
 Q00651

 Gene ID
 16401

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

CD49d is a 150 kD glycoprotein, also known as α4 integrin or VLA-4 α chain. It is a member of the integrin family, expressed on T and B cells, monocytes, eosinophils, basophils, mast cells, thymocytes, NK cells, and dendritic cells. CD49d is a heterodimer expressed with either of two β chains, β 1 (CD29) or β 7, to form the VLA-4 (integrin $\alpha 4\beta 1)$ or LPAM-1 (integrin $\alpha 4\beta 7)$ complexes. CD49d plays a critical role in adhesion and T cell costimulation. The primary ligands for CD49d are VCAM-1, MAdCAM-1, and fibronectin.

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web:www.elabscience.com