

Human PLAU/uPA Antibody Pair Set

Catalog No.	E-KAB-0186	Applications	ELISA
Synonyms	UPA, u-PA, ATF, URK, UP-A, Urokinase, Abbokinase		

Kit components & Storage

Title	Specifications	Storage
Human PLAU/uPA Capture Antibody	1 vial, 100 µg	Store at -20°C for one year. Avoid freeze / thaw cycles.
Human PLAU/uPA Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20°C for one year. Avoid freeze / thaw cycles.


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0186)	
		Human PLAU/uPA Capture Antibody	Human PLAU/uPA Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human PLAU/uPA protein	Recombinant Human PLAU/uPA protein
	Swissprot	P00749	
Product details	Reactivity	Human	Human
	Host	Mouse	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Human PLAU/uPA in ELISAs.	

Applications

Human PLAU/uPA Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4µg/mL	Human PLAU/uPA Capture Antibody	 <p>The graph displays a standard curve for the Human PLAU/uPA Sandwich ELISA Assay. The y-axis represents Optical Density, ranging from 0.01 to 10 on a logarithmic scale. The x-axis represents Human PLAU/uPA concentration in pg/mL, ranging from 10 to 100,000 on a logarithmic scale. Six data points are plotted, showing a clear positive linear relationship on the log-log scale, indicating a wide dynamic range for the assay.</p>
ELISA Detection	1:1000-1:10000	Human PLAU/uPA Detection Antibody (Biotin)	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

This gene encodes a serine protease involved in degradation of the extracellular matrix and possibly tumor cell migration and proliferation. A specific polymorphism in this gene may be associated with late-onset Alzheimer's disease and also with decreased affinity for fibrin-binding. This protein converts plasminogen to plasmin by specific cleavage of an Arg-Val bond in plasminogen. Plasmin in turn cleaves this protein at a Lys-Ile bond to form a two-chain derivative in which a single disulfide bond connects the amino-terminal A-chain to the catalytically active, carboxy-terminal B-chain. This two-chain derivative is also called HMW-uPA (high molecular weight uPA). HMW-uPA can be further processed into LMW-uPA (low molecular weight uPA) by cleavage of chain A into a short chain A (A1) and an amino-terminal fragment. LMW-uPA is proteolytically active but does not bind to the uPA receptor. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.