# N/H/K-Ras Polyclonal Antibody

catalog number: E-AB-70313



Note: Centrifuge before opening to ensure complete recovery of vial contents.

## Description

**Reactivity** Mouse; Rat

Immunogen KLH conjugated Synthetic peptide corresponding to Mouse K-Ras

Host Rabbit
Isotype IgG

Purification Affinity purification
Conjugation Unconjugated


buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 1% protein

protectant and 50% glycerol.

**Applications** Recommended Dilution

**WB** 1:500-1:2000

#### Data



Western Blot analysis of various samples using N/H/K-Ras

Polyclonal Antibody at dilution of 1:800.

Observed-MV:21 kDa Calculated-MV:21 kDa

### **Preparation & Storage**

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

## Background

NRAS an N-ras oncogene encoding a membrane protein that shuttles between the Golgi apparatus and the plasma membrane. This shuttling is regulated through palmitoylation and depalmitoylation by the ZDHHC9-GOLGA7 complex. The encoded protein, which has intrinsic GTPase activity, is activated by a guanine nucleotide-exchange factor and inactivated by a GTPase activating protein. Mutations in this gene have been associated with somatic rectal cancer, follicular thyroid cancer, autoimmune lymphoproliferative syndrome, Noonan syndrome, and juvenile myelomonocytic leukemia. HRAS belongs to the Ras oncogene family, whose members are related to the transforming genes of mammalian sarcoma retroviruses. The products encoded by these genes function in signal transduction pathways. These proteins can bind GTP and GDP, and they have intrinsic GTPase activity. KRAS, a Kirsten ras oncogene homolog from the mammalian ras gene family, encodes a protein that is a member of the small GTPase superfamily. A single amino acid substitution is responsible for an activating mutation. The transforming protein that results is implicated in various malignancies, including lung adenocarcinoma, mucinous adenoma, ductal carcinoma of the pancreas and colorectal carcinoma.

## For Research Use Only