

Myc-Tag Monoclonal Antibody

catalog number: **E-AB-48023**

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description


Reactivity	All
Immunogen	Synthetic peptide corresponding to Myc tag conjugated to keyhole limpet haemocyanin.
Host	Mouse
Isotype	IgG2a
Clone	3E9
Purification	Protein A/G Purification
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

Recommended Dilution

WB	1:2000-4000
FCM	0.2µg


Data



Western blot with Anti Myc-Tag Monoclonal Antibody at dilution of 1:4000. Lane 1: 293T Transfected with Myc-Tag cell lysate, Lane 2: 293T cell lysate.

Observed-MW: 28 kDa

Calculated-MW: 28 kDa



1x10⁶ CHO cells Transfected with a MYC plasmid were stained with 0.2µg Anti-MYC tag mAb(E-AB-48023) and AF647 conjugated Goat Anti-mouse IgG(H+L); Isotype Control stained with 0.2µg mouse IgG and AF647 conjugated Goat Anti-mouse IgG (H+L)

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

The MYC tag, derived from the c-MYC protein, is a popular epitope tag for detecting the expression of recombinant proteins in yeast, bacteria, insect, and mammalian cell systems. It can be used for affinity chromatography, then used to separate recombinant, overexpressed protein from wild type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V2.8