

Recombinant Human CHI3L2/YKL-39 Protein (His Tag)

Catalog Number: PKSH033718

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human CHI3L2;YKL-39 protein Tyr27-Leu390, with an C-terminal His
Mol_Mass	41.9 kDa
Accession	AAH11460.1
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, 5% Trehalose, 5% Mannitol, 0.01% Tween 80, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Chitinase 3-Like Protein 2 (CHI3L2) is a 39 kDa secreted glycoprotein which belongs to the glycosyl hydrolase 18 family and the most closely related to human cartilage glycoprotein 39, which promotes the growth of human synovial cells as well as skin and fetal lung fibroblasts. Highest expression of CHI3L2 is in chondrocytes, followed by synoviocytes, lung and heart. It is not detected in spleen, pancreas, and liver. CHI3L2 may also be expressed in developing brain and placenta. Increased levels of CHI3L2 have been demonstrated in synovial fluids of patients with rheumatoid or osteoarthritis as well as in some other pathologies and in malignant tumors, particularly in glioblastomas. CHI3L2 may bind glycan structure with high affinity, but not heparin. It has no chitotriosidase activity, but is likely to bind some type of glycan.