Recombinant Human Cystatin C/CST3 Protein (Human Cells, His Tag)

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Catalog Number: PKSH032323



Description **Species** Human 14.4 kDa Mol Mass Accession P01034 **Bio-activity** Not validated for activity **Properties** > 95 % as determined by reducing SDS-PAGE. Purity Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method. Store at $< -20^{\circ}$ C, stable for 6 months. Please minimize freeze-thaw cycles. Storage This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel Shipping packs. Upon receipt, store it immediately at $< -20^{\circ}$ C. Formulation Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 500mM NaCl, 0.01% Tween 80, 1mM EDTA, 50% Glycerol, pH 8.0. Reconstitution Not Applicable Data N 417

kDa	MK	R
120 90		
60		
40	-	
30	-	
20	-	
14		

> 95 % as determined by reducing SDS-PAGE.

Background

Cystatin C is a member of family 2 of the cystatin superfamily. It is ubiquitous in human tissues and body fluids and mainly used as a biomarker of kidney function. Cystatin C inhibits many cysteine proteases such as papain and Cathepsins B; H; K; L and S. As an inhibitor of cysteine proteinases; Cystatin C is thought to serve an important physiological role as a local regulator of this enzyme activity. Recently; it has been studied for its role in predicting newonset or deteriorating cardiovascular disease. It also seems to play a role in brain disorders involving amyloid (a specific type of protein deposition); such as Alzheimer's disease.

For Research Use Only