

A Reliable Research Partner in Life Science and Medicine

PE Anti-Human CD10 Antibody[HI10a]

Catalog Number: E-AB-F1141D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Mouse Host

Isotype Mouse IgG1, ĸ

Clone No. HI10a

PE Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792D] Isotype Control

Conjugation

Conjugation Information PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



Human peripheral blood leucocytes are stained with PE Anti-Human CD10 Antibody (Left). Leucocytes are stained with PE Mouse IgG1, κ Isotype Control (Right).

Preparation & Storage

Keep as concentrated solution. **Storage**

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names Atriopeptidase:CALLA:CD10:Enkephalinase:NEP:Neprilysin:Neutral endopeptidase:

SFE;Skin fibroblast elastase

Uniprot ID P08473 Gene ID 4311

For Research Use Only

Fax: 1-832-243-6017 Tel: 1-832-243-6086 Toll-free: 1-888-852-8623 Email:techsupport@elabscience.com

Web:www.elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

CD10 is a 100 kD neutral endopeptidase and a member of the metalloprotease family. It is a type II transmembrane protein also known as common acute lymphoblastic leukemia antigen (CALLA), enkephalinase, and neprilysin. CD10 is expressed on B cell precursors, T cell precursors, and neutrophils. CD10 is involved in B cell development and has been shown to bind opioid enkephalins, bradykinin, angiotensins I and II, and other biologically active peptides.

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web:www.elabscience.com