

TTLL12 Polyclonal Antibody

catalog number: E-AB-92358

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|---------------------|--|
| Reactivity | Human |
| Immunogen | Recombinant fusion protein of human TTLL12 |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

Applications

| | |
|-----------|--------------|
| WB | 1:500-1:2000 |
|-----------|--------------|

Data



Western blot analysis of extracts of various cell lines using
TTLL12 Polyclonal Antibody at 1:1000 dilution.

Observed-MW:Refer to figures

Calculated-MW:74 kDa

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

Background

Negatively regulates post-translational modifications of tubulin, including detyrosination of the C-terminus and polyglutamylation of glutamate residues. Also, indirectly promotes histone H4 trimethylation at 'Lys-20' (H4K20me3). Probably by controlling tubulin and/or histone H4 post-translational modifications, plays a role in mitosis and in maintaining chromosome number stability. During RNA virus-mediated infection, acts as a negative regulator of the DDX58/RIG-I pathway by preventing MAVS binding to TBK1 and IKKε.

For Research Use Only