TTLL12 Polyclonal Antibody

catalog number: **E-AB-92358**


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	Human
Immunogen	Recombinant fusion protein of human TTLL12
Host	Rabbit
Isotype	Rabbit IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilution
WB	1:500-1:2000

Data



Western blot analysis of extracts of various cell lines using TTLL12 Polyclonal Antibody at 1:1000 dilution.

Observed-MW:Refer to figures

Calculated-MW:74 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

Negatively regulates post-translational modifications of tubulin, including detyrosination of the C-terminus and polyglutamylation of glutamate residues. Also, indirectly promotes histone H4 trimethylation at 'Lys-20' (H4K20me3. Probably by controlling tubulin and/or histone H4 post-translational modifications, plays a role in mitosis and in maintaining chromosome number stability. During RNA virus-mediated infection, acts as a negative regulator of the DDX58/RIG-I pathway by preventing MAVS binding to TBK1 and IKBKE.