Elabscience®

COX11 Polyclonal Antibody

catalog number: E-AB-14015

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Reactivity	Human;Mouse	
Immunogen	Recombinant protein of human COX11	
Host	Rabbit	
Is otype	IgG	
Purification	Affinity purification	
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	
Applications	Recommended Dilution	
WB	1:1000-1:5000	
IHC	1:25-1:100	
Data		
9- 2- 3- 3- 3- 9-	brain tissue using COX11	Immunohistochemistry of paraffin-embedded Human
Western Blot analysis of Mouse brain tissue using COX11 Polyclonal Antibody at dilution of 1:600		stomach cancer using COX11 Polyclonal Antibody at
Calculated-MW:31 kDa		dilution of 1:30
Preparation & Storage	, w i nDa	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.	
Shipping	The product is shipped with ice pack,upon receipt, store it immediately at the	
Smblving	temperature recommended.	

Background

Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation and assembly of the complex This nuclear gene encodes a protein which is not a structural subunit, but may be a heme A biosynthetic enzyme involved in COX formation, according to the yeast mutant studies. However, the studies in Rhodobacter sphaeroides suggest that this gene is not required for heme A biosynthesis, but required for stable formation of the Cu(B) and magnesium centers of COX. This human protein is predicted to contain a transmembrane domain localized in the mitochondrial inner membrane. Multiple transcript variants encoding different isoforms have been found for this gene. A related pseudogene has been found on chromosome 6.

For Research Use Only