

## Recombinant Rat ASAM Protein (His Tag)

**Catalog Number:** PKSR030312

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Rat
<b>Source</b>	HEK293 Cells-derived Rat ASAM protein Met1-Met232, with an C-terminal His
<b>Calculated MW</b>	25.6 kDa
<b>Observed MW</b>	37 kDa
<b>Accession</b>	Q8K1G0
<b>Bio-activity</b>	Measured by the ability of the immobilized protein to support the adhesion of MS-1 cells. When $5 \times 10^4$ cells/well are added to Recombinant Rat ASAM coated plates (12.5 µg/mL with 100 µL/well), 35-70% cells will adhere after 1 hour incubation at 37°C.

### Properties

<b>Purity</b>	> 85 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 85 % as determined by reducing SDS-PAGE.

### Background

Adipocyte-specific adhesion molecule (ASAM), also known as ACAM and CLMP, is a type I transmembrane protein and a member of the CTX (cortical thymocyte marker in Xenopus) family within the immunoglobulin superfamily. ASAM protein is highly expressed in the small intestine and placenta, and is found at intermediate levels in the heart, skeletal muscle, colon, spleen, kidney, and lung, and appears in low levels in the liver and peripheral blood leukocytes as well. ASAM is a transmembrane component of tight junctions in epithelial cells that can mediate cell aggregation and regulate transepithelial resistance across polarized epithelial cells. In addition, its expression is strongly correlated with white adipose tissue (WAT) mass of human and rodents with obesity.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017