

Recombinant Human S100A4 Protein (Fc Tag)

Catalog Number: PKSH031787

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human S100A4 protein Met 1-Lys 101, with an N-terminal hFc
Calculated MW	38.4 kDa
Observed MW	40 kDa
Accession	NP_002952.1
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

S100A4, also known as metastasis-associated protein Mts1, belongs to the family of small calcium-binding S100 proteins containing two EF-hand calcium-binding motifs. In humans at least 20 S100 family members that are distributed tissue specifically have been identified, and are involved in a number of cellular processes as transducers of calcium signal. S100A4 is a symmetric homodimer, and undergoes a relatively large conformational change upon the typical EF-hand binding calcium, which is necessary for S100A4 to interact with its protein targets and generate biological effects. It can bind the already known targets p53, F-actin, liprin β , myosin heavy chain II, and prevent their phosphorylation and multimerization. It has been demonstrated that S100A4 is directly involved in tumor metastasis including cell motility, invasion, apoptosis, angiogenesis and differentiation, and appears to be a metastasis factor and a molecular marker for clinical prognosis. Multiple alternatively spliced variants encoding the same protein have been identified.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017