
Recombinant Human AIF1 Protein(Trx Tag)

Catalog Number: PDEH100611

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	E.coli-derived Human AIF1 protein Met1-Pro147, with an N-terminal Trx
Calculated MW	36 kDa
Observed MW	36 kDa
Accession	P55008-1
Bio-activity	Not validated for activity
Properties	
Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with 5% Trehalose and 5%
	Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of
	0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human AIF1 proteins, 2µg/lane of Recombinant Human AIF1 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 36 KD

Background

AIF1, also known as IBA1, is an actin-binding protein. AIF1 is expressed selectively in human macrophage-like cell lines, and in a subset of CD68(+) macrophages in the interstitial and perivascular spaces of human heart allografts. It is expressed in macrophages and neutrophils. AIF1 enhances membrane ruffling and RAC activation. AIF1 enhances the actin-bundling activity of LCP1. It also enhances lymphocyte migration. AIF1 may play a role in macrophage activation and function. It binds calcium and plays a role in RAC signaling and in phagocytosis. It promotes the proliferation of vascular smooth muscle cells and of T-lymphocytes.

For Research Use Only Toll-free: 1-888-852-8623