

Biotin Anti-Mouse CD80 Antibody[16-10A1]

Catalog Number: E-AB-F0992B

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	Mouse
Host	Armenian Hamster
Isotype	Armenian Hamster IgG
Clone No.	16-10A1
Isotype Control	Biotin Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09853B]
Conjugation	Biotin
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \mu\text{g}$ per 10^6 cells in 100 μL volume or 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
------------	---

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	Activation B7-1 antigen;B7;Cd80;T-lymphocyte activation antigen CD80
Uniprot ID	Q00609
Gene ID	12519
Background	CD80 is a 60 kD highly glycosylated protein. It is a member of the Ig superfamily and is also known as B7-1, B7, and Ly-53. CD80 is constitutively expressed on dendritic cells and monocytes/macrophages, and inducibly expressed on activated B and T cells. The ligation of CD28 on T cells with CD80 and CD86 (B7-2) on antigen presenting cells (such as dendritic cells, macrophages, and B cells) elicits co-stimulation of T cells resulting in enhanced cell activation, proliferation, and cytokine production. CD80 appears to be expressed later in the immune response than CD86. CD80 can also bind to CD152, also known as CTLA-4, to deliver an inhibitory signal to T cells.

For Research Use Only