



A Reliable Research Partner in Life Science and Medicine

# **Human PGF Antibody Pair Set**

Catalog No. E-KAB-0221 Applications ELISA

Synonyms PLGF, PGFL, PIGF-2, SHGC-10760, D12S1900, PIGF

## **Kit components & Storage**

Title	Specifications	Storage
Human PGF Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Human PGF Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20℃ for one year.
		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0221)		
		Human PGF Capture Antibody	Human PGF Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human PGF protein	Recombinant Human PGF protein	
Information	Swissprot	P49763		
Product details	Reactivity	Human	Human	
	Host	Mouse	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Protein A or G	Antigen Affinity	
	Specificity	Detects Human PGF in ELISAs.		

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>



A Reliable Research Partner in Life Science and Medicine

## **Applications**

Human PGF Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Human PGF Capture Antibody	
Capture			Aist
ELISA Detection	1:1000-1:10000	Human PGF Detection Antibody (Biotin)	0.01 100 1000 10000 Human PGF concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

### **Background**

Placental growth factor is a homodimeric glycoprotein, 46-50 kDa in size, belonging to the vascular endothelial growth factor (VEGF) sub-family. It plays an important role in pathological VEGF-driven angiogenesis. PLGF is expressed not only in placental cells but also in colon and mammary carcinomas. It acts as a primary inflammatory instigator of atherosclerotic plaque instability and thus may be useful as a risk-predicting biomarker in patients with acute coronary syndromes (ACS).

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>