Purified Anti-Mouse CD69 Antibody[H1.2F3]

catalog number: **E-AB-F1187A**


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	Mouse
Immunogen	Recombinant Human CD69 protein
Host	Armenian Hamster
Isotype	Armenian Hamster IgG
Clone	H1.2F3
Purification	>98%, Protein A/G purified
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications	Recommended Dilution
FCM	≤ 0.2 μg per million cells in 100 μL volume

Data



HEK293T cells transfected with pcDNA3.1 plasmid encoding Mouse CD69 gene were stained with 0.2 μg Purified Anti-Mouse CD69 Antibody[H1.2F3] (Right) and 0.2 μg Armenian Hamster IgG, κ Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Armenian Hamster IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

CD69 is a 60 kD type II membrane protein composed of a 27/33 kD disulfide-linked homodimer, also known as Very Early Activation Antigen (VEA), AIM, EA1, MLR3, and gp34/28. It is expressed on a subset of thymocytes and platelet s. CD69 is rapidly induced on activated T and B cells, neutrophils, and NK cells. It is a C-type lectin, closely related to the NKR-P1 and Ly-49 NK cell activation molecules. CD69 is involved in the early events of cell activation and thymocyte positive selection.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com