



A Reliable Research Partner in Life Science and Medicine

## APC Anti-Human CD72 Antibody[3F3]

Catalog Number: AN00325E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Mouse Host

Isotype Mouse IgG2b, ĸ

Clone No. 3F3

APC Mouse IgG2b, k Isotype Control[MPC-11] [Product E-AB-F09812E] Isotype Control

Conjugation

**Conjugation Information** APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

**Applications** Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount **FCM** 

> of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

**Preparation & Storage** 

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

**Shipping** Ice bag

**Antigen Information** 

Lyb-2;Ly-19.2;Ly-32.2 **Alternate Names** 

**Uniprot ID** P21854 Gene ID 971

**Background** CD72 is a 39-43 kD type II membrane glycoprotein. It is a disulfide-linked homodimer

> belonging to C-type lectin family. CD72 is a pan-B cell marker expressed on pre-pre-B cells throughout B cell differentiation with the exception of plasma cells. It is also expressed on follicular dendritic cells, splenic red pulp macrophages (but not on peripheral blood monocytes), and liver Kupffer cells. CD72, a negative coreceptor of B cells, contains immunoreceptor tyrosine-based inhibitory motifs in the cytoplasmic domain which has been shown to recruit the tyrosine phosphatase SHP-1. Ligation of

CD72 with its ligand regulates CD72 tyrosine dephosphorylation and SHP-1

dissociation to promote B cell activation and proliferation. CD100 and CD5 have been shown to be CD72 ligands. The CD100-CD72 interaction plays a role in maintenance

of B cell homeostasis.

Fax: 1-832-243-6017 Tel: 1-832-243-6086 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Email:techsupport@elabscience.com