Purified Anti-Rat CD45 Antibody[OX-1]

catalog number: E-AB-F1227A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Reactivity	Rat
Immunogen	Recombinant Rat CD45 protein
Host	Mouse
lsotype	Mouse IgG1, ĸ
Clone	OX-1
Purification	>98%, Protein A/G purified
Buffer	PBS, pH 7.2. Contains 0.05% Proclin300.
Applications	Recommended Dilution
FCM	$2 \ \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$
Data	
Commt (%)	
Rat splenocytes were sta CD45 Antibody[OX-1] Isotype Control (Left)	$\int_{1}^{2} \int_{1}^{10^3} \int_{10^4}^{10^5} \int_{10^6}^{10^6}$ $\int_{2}^{10^3} \int_{10^4}^{10^5} \int_{10^6}^{10^6}$ $\int_{10^4}^{10^5} \int_{10^6}^{10^6}$ $\int_{10^6}^{10^6} \int_{10^6}^{10^6}$ $\int_{10^6}^{10^6} \int_{10^6}^{10^6} \int_{10^6}^{10^6}$ $\int_{10^6}^{10^6} \int_{10^6}^{10^6} \int_{10^6}^{10^6}$ $\int_{10^6}^{10^6} \int_{10^6}^{10^6} \int_{10^6}^{10^6} \int_{10^6}^{10^6}$ $\int_{10^6}^{10^6} \int_{10^6}^{10^6} \int_{$
Rat splenocytes were sta CD45 Antibody[OX-1] Isotype Control (Left) conjugated Goat Anti-	CD15 Alexa Fluor® 647+1 Lined with 0.2 μg Purified Anti-Rat (Right) and 0.2 μg Mouse IgG1, κ , followed by Alexa Fluor® 647-
Rat splenocytes were sta CD45 Antibody[OX-1] Isotype Control (Left) conjugated Goat Anti- Preparation & Storage	CD15 Alexa Fluor® 647+1 Lined with 0.2 μg Purified Anti-Rat (Right) and 0.2 μg Mouse IgG1, κ , followed by Alexa Fluor® 647-
Rat splenocytes were sta CD45 Antibody[OX-1] Isotype Control (Left)	CDU5 Alexa Fluor® 647+1 Lined with 0.2 μg Purified Anti-Rat (Right) and 0.2 μg Mouse IgG1, κ , followed by Alexa Fluor® 647- Mouse IgG Secondary Antibody. Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

extracellular domain and glycosylation. CD45 is expressed on all hematopoietic cells except erythrocytes and platelets. CD45RA is one of the CD45 isoforms with a molecular weight of 200-220 kD. It is expressed almost exclusively on B cells. CD45 functions in signal transduction through T and B cell antigen receptors. CD45 has been shown to interact with various proteins, including galectin-1, CD2, CD3, and CD4.