BDH2 Polyclonal Antibody

catalog number: E-AB-18535



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat **Immunogen** Full length fusion protein

Host Rabbit Isotype IgG

Purification Antigen affinity purification


Conjugation Unconjugated

buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilution

WB 1:500-1:2000 **IHC** 1:50-1:300

Data





Western blot analysis of Mouse kidney tissue and A172 cell Immunohistochemistry of paraffin-embedded Human liver lysates using BDH2 Polyclonal Antibody at dilution of 1:600 cancer tissue using BDH2 Polyclonal Antibody at dilution of 1:650(×200)

serveu-iviv. Reier to figu

Calculated-MV:27 kDa

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

Dehydrogenase that mediates the formation of 2,5-dihydroxybenzoic acid (2,5-DHBA), a siderophore that shares structural similarities with bacterial enterobactin and associates with LCN2, thereby playing a key role in iron homeostasis and transport. Also acts as a 3-hydroxybutyrate dehydrogenase (By similarity).

For Research Use Only