

EIF2 alpha Polyclonal Antibody

catalog number: E-AB-31295

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|---------------------|---|
| Reactivity | Human;Mouse;Rat;Monkey |
| Immunogen | Synthesized peptide derived from human eIF2 α around the non-phosphorylation site of Ser51. |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Conjugation | Unconjugated |
| Buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein protectant and 50% glycerol. |

Applications

Recommended Dilution

| | |
|------------|--------------|
| WB | 1:500-1:2000 |
| IHC | 1:100-1:300 |
| IF | 1:50-1:200 |


Data




Western Blot analysis of various cells using EIF2 alpha Polyclonal Antibody at dilution of 1:1000.

Observed-MW:38 kDa

Calculated-MW:36 kDa



Immunofluorescence analysis of Rat kidney tissue using EIF2 alpha Polyclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using EIF2 alpha Polyclonal Antibody at dilution of 1:200.

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

Background

For Research Use Only

Functions in the early steps of protein synthesis of a small number of specific mRNAs. Acts by directing the binding of methionyl-tRNA_i to 40S ribosomal subunits. In contrast to the eIF-2 complex, it binds methionyl-tRNA_i to 40 S subunits in a codon-dependent manner, whereas the eIF-2 complex binds methionyl-tRNA_i to 40 S subunits in a GTP-dependent manner. May act by impinging the expression of specific proteins.