

HDAC2 Polyclonal Antibody

catalog number: E-AB-31644

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|---------------------|---|
| Reactivity | Human;Mouse;Rat;Monkey |
| Immunogen | Synthesized peptide derived from human HDAC2 around the non-phosphorylation site of Ser394. |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein protectant and 50% glycerol. |

Applications

Recommended Dilution

| | |
|------------|--------------|
| WB | 1:500-1:2000 |
| IHC | 1:100-1:300 |

Data



Western Blot analysis of Jurkat, 3T3 cells using HDAC2 Polyclonal Antibody at dilution of 1:2000.

Observed-MW:55 kDa

Calculated-MW:55 kDa

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

Background

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Forms transcriptional repressor complexes by associating with MAD, SIN3, YY1 and N-COR. Interacts in the late S-phase of DNA-replication with DNMT1 in the other transcriptional repressor complex composed of DNMT1, DMAP1, PCNA, CAF1. Deacetylates TSHZ3 and regulates its transcriptional repressor activity.

For Research Use Only