



A Reliable Research Partner in Life Science and Medicine

Purified Anti-Mouse CD18 Antibody[M18/2]

catalog number: E-AB-F1113A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse

Immunogen Recombinant Mouse CD18 protein

Host Rat

Isotype Rat IgG2a, κ
Clone M18/2

Purification >98%, Protein A/G purified


Buffer Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer.

Dialyze to completely remove the stabilizer prior to labeling.

Applications Recommended Dilution

FCM 2 μ g/mL(0.5×10⁶-1×10⁶ cells)

Data



C57/BL6 Mouse splenocytes were stained with 0.2 μg Purified Anti-Mouse CD18 Antibody[M18/2] (Right) and 0.2 μg Rat IgG2a, κ Isotype Control (Left), followed by Elab

Fluor[®] 647-conjugated Goat Anti-Rat IgG Secondary Antibody.

Preparation & Storage

Storage Storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

CD18 is a 95 kD protein, also known as integrin β 2 subunit. It is expressed on all leukocytes. CD18, in association with integrin α chain CD11a, CD11b, and CD11c forms LFA-1, Mac-1, and α X β 2, respectively, and plays an important role in leukocytes adhesion. The CD18 integrin complexes bind ICAM-1 (CD54), ICAM-2 (CD102), ICAM-3 (CD50), iC3b, and fibrinogen. The M18/2 antibody is able to block tumor cell metastasis and enhance cell adhesion.

For Research Use Only