

Purified Anti-Mouse CD18 Antibody[M18/2]

catalog number: E-AB-F1113A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Host	Rat
Isotype	Rat IgG2a, κ
Clone	M18/2
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

Recommended Dilution

FCM	2 μ g/mL(1×10^5 - 5×10^5 cells)
------------	--

Data



C57/BL6 Mouse bone marrow cells were stained with 0.2 μ g Purified Anti-Mouse CD18 Antibody[M18/2] (Right) and 0.2 μ g Rat IgG2a, κ Isotype Control (Left), followed by PE-conjugated Goat Anti-Rat IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

CD18 is a 95 kD protein, also known as integrin $\beta 2$ subunit. It is expressed on all leukocytes. CD18, in association with integrin α chain CD11a, CD11b, and CD11c forms LFA-1, Mac-1, and $\alpha X\beta 2$, respectively, and plays an important role in leukocytes adhesion. The CD18 integrin complexes bind ICAM-1 (CD54), ICAM-2 (CD102), ICAM-3 (CD50), iC3b, and fibrinogen.

For Research Use Only